



# Pest e-alerts



---

Entomology and Plant Pathology, Oklahoma State University  
127 Noble Research Center, Stillwater, OK 74078  
405.744.5527

---

Vol. 18, No. 2

<http://entopl.okstate.edu/pddl/pdidl>

2/8/2019

---

## Wheat Disease Update – 8 February 2019

Bob Hunger, Extension Wheat Pathologist  
Department of Entomology & Plant Pathology  
Oklahoma State University - 127 Noble Research Center  
405-744-9958

This is just a brief update from a discussion I had today with Gary Strickland (Extension Educator, Jackson County in southwestern OK). He indicated the majority of wheat in southwestern OK is quite small due to late (mostly November) planting. He has seen some spotty plant death in some fields that appears to be common root rot as there is significant darkening of sub-crown internodes (Figure 1). In addition, he has observed leaf rust on the lowest leaves close to the ground in a number of the varieties at the variety trial located near Altus. He hasn't seen rust at any other location, but did find it on wheat at the Altus variety trial, which he indicated was some of the bigger wheat in southwestern OK.

Figure 1. Common root rot on wheat. Note the darkened sub-crown internodes (left and center photo) compared to a creamish-white (healthy) sub-crown internode on the right.



**Reports from other states:** Leaf rust appears to be increasing across southern and central Texas as indicated by Dr. Clark Neely (Small Grains and Cool-season Oilseed Extension Specialist for Texas A&M AgriLife Extension). Dr. Neely reported finding leaf rust in mid-January near College Station, TX, but adds to this by reporting, “I heard my first report of fairly widespread leaf rust in producer fields from a crop consultant in the Temple, TX area last week (Central Texas). It was reportedly observed on ‘WB Cedar’ which is known to be susceptible to leaf rust and was seen in multiple fields at varying levels of infection. More cloudy, rainy weather is forecasted for the next 10 days with highs fluctuating from mid 40’s to mid 70’s.”

So, taking this report along with Gary Strickland’s observation indicates that as temperature increases producers need to be watching for rust (both leaf and stripe although only leaf rust has been found so far) in southern OK. Early infection by stripe rust does not always produce pustules in stripes as seen with later infections. It is important to be aware of this as it is easy to confuse leaf rust and stripe rust juvenile infections if one is looking for pustules in stripes. Typically the color of the spores in the pustules is different with stripe rust being more yellowish/golden orange versus a darker brownish red for leaf rust (Figure 2).

Figure 2. Stripe rust early (juvenile) infection (left photo; note how pustules are not in a stripe); stripe rust adult infection (center – Dr. Mary Burrows; Montana State University); stripe rust and leaf rust on the same leaf (right photo – Dr. Stephen Harrison; Louisiana State University). Note the darker brown (rusty) color of leaf rust pustules compared to the more yellowish golden orange color of stripe rust pustules.



#### Disease and Insect Diagnostic Laboratory

The pesticide information presented in this publication was current with federal and state regulations at the time of printing. The user is responsible for determining that the intended use is consistent with the label of the product being used. Use pesticides safely. Read and follow label directions. The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: [eeo@okstate.edu](mailto:eeo@okstate.edu) has been designated to handle inquiries regarding non-discrimination policies; Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU’s Title IX Coordinator 405-744-9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources.