



Pest e-alerts



*Entomology and Plant Pathology, Oklahoma State University
127 Noble Research Center, Stillwater, OK 74078
405.744.5527*

Vol. 18, No. 16

<http://entopl.okstate.edu/pddl/pddl>

5/3/19

Wheat Disease Update – 3 May 2019

Bob Hunger, Extension Wheat Pathologist
Department of Entomology & Plant Pathology
Oklahoma State University - 127 Noble Research Center
405-744-9958

Reports and observations of foliar diseases increased this past week. Around Stillwater, wheat ranges from heads nearly or completely emerged to full kernel (watery). On this wheat, higher levels of leaf and stripe rust were observed in various trials and nurseries (Figure 1). There seems to be more chlorosis/yellowing associated with these foliar diseases than in most years. Septoria also is widespread around Stillwater as it is around much of the state. At a field day on 1-May near Apache, OK in Caddo County (central OK), low levels of active stripe rust and leaf rust were observed on susceptible varieties, but by far the most prevalent disease was septoria (Figure 2). Typically the leaf spot diseases (primarily septoria and tan spot) do not move much above the lower leaves, and by this time in May drier weather and higher temperature prevent these diseases from moving up the canopy. However, this year the extended and consistent cloudy, cool and wet conditions have obviously favored septoria to the point that septoria could be found on the leaf just under the flag (but not near as severe as on the lower and mid leaves). Comparing the top two photos in Figure 2 with the bottom photo shows the benefit from applying a fungicide sufficiently early to help manage septoria as the plot in the bottom photo received a fungicide application about 3 weeks earlier (12-April). For more information on applying fungicides and their relative effectiveness in managing foliar diseases, see OCES Current Report (CR-7668) that can be found at:

<http://pods.dasnr.okstate.edu/docushare/dsweb/Get/Document-4987/CR-7668web2019.pdf>

Figure 1. General yellowing of upper foliage (left photo) due primarily to leaf rust infection (right photo) on wheat near Stillwater, OK.



Figure 2. Septoria on mid-foliage of wheat at Apache, OK



Figure 2 (Continue) a close up of an individual leaf (upper photo). Not the small, black “pepper spots” on the leaf (hard to see – very small), which are the spore containing structures of the fungus that causes this disease. The bottom photo shows the lower canopy of a split plot of this wheat that was sprayed with a foliar fungicide three weeks earlier.




Another symptom that is becoming more apparent across Oklahoma is physiological leaf spot (PLS), which we tend to see in years with extended cool and wet weather. Figure 3 shows PLS as observed this past week at Apache along with clearer photos of PLS taken in past years. This type of spotting often is attributed to rust infection as when you look “through” the leaf the spots appear to be about the same as a leaf rust pustule. However, in past years I have taken such plants to the greenhouse and rust never developed. I’ve never seen any papers indicating how much harm PLS can inflict, nor any suggestions to reduce or control its occurrence. In most years and places I have seen it occur, its effect on yield appears to be minimal. However, typically I’ve seen it occur in early to mid-April rather than in early to mid-May.

Figure 3. Physiological leaf spotting (PLS) on wheat. The photo is on wheat at Apache, OK on 1-May.



Figure 3 (Continue) The top and bottom photos are much better (more clear) photos of PLS taken in 2012 (middle) and 2008 (bottom).



So to summarize, wheat foliar diseases are increasing across Oklahoma with septoria being the primary leaf spot disease that is being reported. Persistent wet and cool weather has facilitated movement of septoria up from the lower level foliage into the mid and even to some extent onto upper leaves. There are a number of effective fungicides for these leaf spot diseases, and although application likely would have been most beneficial 7-10 days ago, protection of the flag leaf is most critical. Stripe rust and leaf rust also are increasing. All of these foliar diseases can be managed with fungicides, but the window to legally apply fungicides onto wheat (especially in southern OK) is closing (or may have closed). Hence, be sure to read the label closely and find a chemical that allows application related to the stage of the wheat. Some fungicides use a growth stage deadline (for example start of flowering or end of flowering), and some use a number of days that must pass between application and harvest (post-application harvest restriction). Also keep in mind the reaction of the variety to these diseases. For example, if your wheat is at or near flowering (heads fully emerged), has good resistance to leaf and stripe rust, and is not showing any or much leaf spotting, applying a fungicide would not be indicated. In contrast, applying a fungicide would be indicated if you have a variety that has good yield potential, is susceptible or moderately susceptible to the rusts and is showing heavy leaf spotting on lower and mid leaves.

Disease and Insect Diagnostic Laboratory

The pesticide information presented in this publication was current with federal and state regulations at the time of printing. The user is responsible for determining that the intended use is consistent with the label of the product being used. Use pesticides safely. Read and follow label directions. The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: eeo@okstate.edu has been designated to handle inquiries regarding non-discrimination policies: Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU's Title IX Coordinator 405-744-9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources.