



Pest e-alerts



Entomology and Plant Pathology, Oklahoma State University
127 Noble Research Center, Stillwater, OK 74078
405.744.5527

Vol. 19, No. 13

<http://entopl.okstate.edu/pddl/pddl>

04/10/2020

Bird Cherry-Oat Aphids and Armyworms in Wheat: To Treat or Not to Treat?

Tom A. Royer, Extension Entomologist: email: tom.royer@okstate.edu, twitter: Tom A. Royer@RoyerTom

It's that time of year! A healthy, good-looking wheat crop provides a "meal" for several wheat pests. I have received scattered reports of some wheat fields infested with bird cherry-oat aphids. Lanie Hale from Wheeler Brothers sent a photo of bird cherry oat aphids in some Blaine County wheat. Bird cherry-oat aphid infestations can be overlooked because they do not produce visible damage until they become very numerous unless they have transmitted the Barley Yellow Dwarf virus. This is an opportune time to check your field bird cherry oat aphid, as well as armyworms!

Bird cherry-oat aphids and a mummy from a wasp parasitoid



My suggestion is to scout the field beforehand to determine if there are GROWING numbers of bird cherry oat aphids that could be or are of concern. Count bird cherry oat aphids on each of 25 randomly selected tillers across a zigzag transect of the field and note mummy activity. If 10-20% of bird-cherry

oat aphids are mummies, and there are numerous lady beetle larvae in the wheat, control may not be warranted.

Unpublished research provided by Dr. Kris Giles (OSU) and Dr. Norm Elliott (USDA-ARS) combined with studies on spring wheat from the Dakotas and Minnesota indicate that 20-40 BCOA per tiller causes 5-9% yield loss before wheat reaches the boot stage. My suggestions: if BCOA numbers average 10-20 per tiller, figure on a 5% loss, if 20-40 per tiller, figure a 7% loss, and if BCOA aphids are more than 40 per tiller, figure a 9% loss.

Estimate APHIDS PER TILLER _____/tiller = Total # aphids _____/25 tillers
 Estimate CROP VALUE \$ _____/acre = Expected yield _____ bushels/acre X \$ _____/bushel
 Calculate CONTROL COSTS \$ _____/acre = Insecticide \$ _____/acre + Application \$ _____/Acre

PREVENTABLE LOSS \$ _____/acre = Crop value \$ _____ X _____ loss from aphids/tiller .

IF PREVENTABLE LOSS IS **GREATER THAN** CONTROL COSTS **TREAT**
 IF PREVENTABLE LOSS IS **LESS THAN** CONTROL COSTS **DON'T TREAT**

Here is a Table of Preventable Loss estimates for bird cherry-oat aphids for expected yields of 30 to 50 bushels per acre, expected wheat prices of \$3.00, \$3.50, and \$4.00 per bushel, and bird cherry-oat aphid numbers of 10-20, 20 to 40, and over 40 per tiller.

Expected Yield (Bushels/Acre)	Expected Price (\$ per bushel)	Crop Value (\$ per acre)	Preventable Loss from BCO Aphids (\$ per acre)		
			10-20 aphids/tiller	20-40 aphids/tiller	More than 40/tiller
30	\$3.50	\$105.00	\$5.25	\$7.35	\$9.45
35	\$3.50	\$122.50	\$6.13	\$8.58	\$11.03
40	\$3.50	\$140.00	\$7.00	\$9.80	\$12.60
45	\$3.50	\$157.50	\$7.88	\$11.30	\$14.18
50	\$3.50	\$175.00	\$8.75	\$12.25	\$15.75
30	\$4.00	\$120.00	\$6.00	\$8.40	\$10.80
35	\$4.00	\$140.00	\$7.00	\$9.80	\$12.60
40	\$4.00	\$160.00	\$8.00	\$11.20	\$14.40
45	\$4.00	\$180.00	\$9.00	\$12.60	\$16.20
50	\$4.00	\$200.00	\$10.00	\$14.00	\$18.00
30	\$4.25	\$127.50	\$6.38	\$8.93	\$11.48
35	\$4.25	\$148.75	\$7.43	\$10.40	\$13.37
40	\$4.25	\$170.00	\$8.50	\$11.90	\$15.30
45	\$4.25	\$191.25	\$9.56	\$13.38	\$17.20
50	\$4.25	\$212.50	\$10.62	\$14.87	\$19.12

This cool, rainy spring weather, while providing excellent growing conditions for wheat, is also feedstuff for “producing” armyworms. Armyworm infestations typically occur in late April through the first two

weeks of May. They damage wheat by feeding on leaves, the awns, and occasionally by clipping the head from developing plants. The head clipping I have noticed over the years is mostly restricted to secondary tillers with very small, green heads that contribute very little to yield.

Armyworm and “head clipping”



Since armyworm infestations occur more frequently around waterways, areas of lush growth, or areas with lodged plants, check them first to determine the size of the infestation. Early signs of an infestation include leaves with ragged margins that have been chewed. You may find “frass” i.e. the excrement from armyworm caterpillars, around the base of wheat stems and clipped heads.


Armyworms: feeding on head, and scouting



Scout for armyworms, at 5 or more locations looking for “curled up worms”. Armyworm caterpillars tend to feed at night, so a good strategy is to bring a flashlight and look at fields after dusk when they are feeding up on the plant stems.

The suggested treatment threshold for armyworms is 4-5 caterpillars per linear foot of row. Generally if wheat is **past** the soft dough stage, control is not warranted unless obvious head clipping can be seen, and caterpillars are present and feeding.

Armyworms: “threshold” numbers of 4-5 per linear foot



If a producer is considering a fungicide application, this might be an opportune time to evaluate your field for bird cherry oat aphid and or armyworms, and IF NEEDED, combine an insecticide with any fungicide application that is needed to control multiple pests. Check [CR-7194](#), "Management of Insect and Mite Pests in Small Grains" for registered insecticides, application rates, and grazing/harvest waiting periods. It can be obtained from any Oklahoma County Extension Office, or found at the OSU Extra Website at <http://pods.dasnr.okstate.edu/docushare/dsweb/Get/Document-2601/CR-7194web2008.pdf>

Co-Editors: Eric Rebek and Justin Talley; Oklahoma Cooperative Extension Service

The pesticide information presented in this publication was current with federal and state regulations at the time of printing. The user is responsible for determining that the intended use is consistent with the label of the product being used. Use pesticides safely. Read and follow label directions. The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: eeo@okstate.edu has been designated to handle inquiries regarding non-discrimination policies; Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU's Title IX Coordinator 405-744- 9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources.