



EXTENSION

Post e-alerts



DEPARTMENT OF
ENTOMOLOGY AND
PLANT PATHOLOGY

*Entomology and Plant Pathology, Oklahoma State University
127 Noble Research Center, Stillwater, OK 74078
405.744.5527*

Vol. 19, No. 22

<http://entopl.okstate.edu/pddl/pdidl>

6/4/2020

Wheat Disease Update – 4 June 2020

Bob Hunger, Extension Wheat Pathologist
Department of Entomology & Plant Pathology
Oklahoma State University - 127 Noble Research Center
405-744-9958

This likely is the last disease update from Oklahoma until preparation starts for planting in the late summer. Harvest is definitely underway across central and southern/southwestern OK, and Josh Bushong (Area Extn Agron Spclst – northwestern and northeastern OK) indicated wheat is quickly turning across northern OK as well. Dry conditions in northwestern OK and the OK panhandle have limited diseases in those parts of OK, and with wheat quickly turning, diseases should not be a factor from here on. The one exception to this would be if head scab was occurring at a high incidence, but to date I have had only one report of just light scab in the northeastern part of Oklahoma.


One report of interest that has come is the occurrence of dark brownish/black to dark bronze-colored heads in mature wheat (Figure 1). Although at a low incidence (Figure 1, left photo) these heads are noticeable and have been observed across multiple varieties by Gary Strickland (Extn Educator; Jackson Cnty; southwestern OK), Dr. Amanda de Oliveira Silva (Ast Professor & Small Grains Extension Agronomist) and Dr. Brett Carver (Regent's Professor/OSU Wheat Geneticist & Breeder), and myself. These heads, which typically are sterile or only have small and shriveled grain, seem to be most common in areas where freeze was most damaging, and likely are related to that environmental cause (i.e., freeze).

Figure 1. Darkened heads at low incidence in mature wheat in a field in southwestern OK. Cause likely is environmental; most likely related to freeze. [photo credit: Gary Strickland; Extn Educator; Jackson Cnty]



I also want to follow-up on the widespread occurrence of splotchy browning and spotting of wheat leaves that was observed across many varieties over Oklahoma this year. I collected and plated numerous leaves from such plants, and the best I can determine is that much of this browning was caused by Septoria leaf blotch (and to a lesser extent, tan spot). Freeze and drought also likely contributed in some cases. Figure 2 shows my reasoning for this. The first four photos are examples of the splotchy browning and spotting of leaves. Some leaves would be almost entirely brown/necrotic. Upon plating and incubating these leaves, I often would not find any pathogens after two days in alternating light dark, which is needed to induce sporulation of the tan spot fungus. However, if I kept these leaves for a longer time (6-10 days), I often would find a high incidence of pycnidia indicative of Septoria leaf blotch forming over the leaves (bottom photo in Figure 1). Oozing from these pycnidia contained spores of the fungus that causes Septoria leaf blotch. Occasionally I also would find sporulation of the fungus that causes tan spot as well as the fungus that causes spot blotch. Also mixed in and often overgrown were many secondary, non-pathogenic fungi that are taking the opportunity to grow on the dead leaf tissue. As a result, I am attributing the widespread leaf browning observed this year to be primarily the result of Septoria leaf blotch.

Figure 2. Various “types” and “degrees” of leaf browning/yellowing/spotting observed on wheat across northern Oklahoma during mid to late May.



So, in summary, the time for wheat diseases in Oklahoma is pretty much at an end for this season in Oklahoma, and I hope everyone has a safe and good harvest!!!

Co-Editors: Eric Rebek and Justin Talley; Oklahoma Cooperative Extension Service

The pesticide information presented in this publication was current with federal and state regulations at the time of printing. The user is responsible for determining that the intended use is consistent with the label of the product being used. Use pesticides safely. Read and follow label directions. The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: eeo@okstate.edu has been designated to handle inquiries regarding non-discrimination policies: Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU's Title IX Coordinator 405-744-9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources.