



Color Patterns in Crossbred Beef Cattle

Megan Rolf
Assistant Professor

Crossbreeding provides commercial cattlemen the opportunity to combine desirable characteristics of two or more breeds (breed complementarity) and increase performance due to hybrid vigor (heterosis). The single strongest argument for crossbreeding is the advantage in fertility and longevity of crossbred cows. With all of the advantages crossbreeding provides, designing an effective mating system should be a top priority for all commercial cattlemen. To plan an effective crossbreeding system, it is helpful to consider any potential detractors that are easy to address. One such detractor exists due to market discrimination against certain breeds and/or colors and lack of uniformity in color. As we increase the number and diversity of breeds involved in crosses, we decrease our ability to maintain complete control of coat color in the offspring unless it was taken into account during breed selection. With that in mind, knowing the basics of coat color inheritance can help producers know what to expect from various breed/composite pairings relative to color pattern.

The purpose of this fact sheet is to provide guidance on how to maintain a uniform color pattern when formulating crossbreeding systems. In some breeds and breed crosses, the color is highly predictable; however, in some other breeds and breed crosses, color is less predictable. A working knowledge of the inheritance of coat color will aid in planning for the color pattern to expect among calves when crossing breeds. This can be very important for evaluating marketing options either at weaning when forming large group lots or on the rail when targeting specific branded beef programs.

Many breeds of beef cattle have a fixed color pattern for that breed because selection has been placed on the color to maintain these characteristics. For example, all Hereford cattle have a red body color with a white face, all Charolais are white, and all Red Poll are red. However, other breeds may

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: <http://osufacts.okstate.edu>

have more than one basic body color such as red or black Limousin or Simmental, and white, red or roan Shorthorn. Still other breeds have multiple colors with more unpredictable inheritance patterns, such as spotting, brindling or stripes in Longhorn and Beefmasters. Some color modifiers under genetic control have been selected against in many breeds (unless they are a feature of color in the breed, such as in Charolais or Hereford) and these features, such as “diluters” and blaze faces, are much less common than in the past, which makes the process of managing color in crossbreeding systems much easier.

Table 1 shows several common breeds of beef cattle and the predominant color pattern that is most commonly associated with each breed. Recently, some breeds with unique color attributes such as spots, blaze faces and diluter genes have been selected against these traits to increase favorable perceptions by terminal buyers. Other breeds that were traditionally red have been selected heavily for black coat color and are listed in Table 1 as both black and red. Introgression of other breeds (specifically Angus) into some of the Continental breeds has altered the traditional color pattern of some of these breeds. Color patterns likely to result from specific crosses are detailed in Table 2.

When you have crossbred cows, predicting color in the offspring can be more difficult, but it helps to understand how color is inherited. All cattle basically possess one of three basic colors: black, red or white. Black is dominant to red, and both black and red are co-dominant with white. One black or red allele with a white allele would result in either a black or red roan animal. In order for an animal to be red or white, they must have two alleles for either red or white,

Table 1. Basic body colors of common cattle breeds.

<i>Black Only</i>	<i>Red Only</i>	<i>Red and Black</i>	<i>White or Cream</i>	<i>Light hair with dark pigmented skin</i>	<i>Spotted</i>	<i>Mixed Colors and/or Roans</i>
Angus	Hereford	Balancer	Charolais	Brahman	Belted Galloway	Beefmaster
Brangus	Lincoln Red	Gelbvieh		Braunvieh	Holstein	Braford
Chiangus	Red Angus	Limousin		Brown Swiss		Longhorn
	Red Poll	Lim-Flex		Jersey		Maine-Anjou
	Santa Gertrudis	Simmental		Nellore		Shorthorn
		Salers				
		SimAngus				

Table 2. Color pattern expected in progeny resulting from the matings of bulls and cows of various colors.

<i>Sire/Dam</i>	<i>Black</i>	<i>Red</i>	<i>White</i>	<i>Light hair with dark skin</i>	<i>Spotted</i>	<i>Mixed Colors</i>
Black	Black (or red if both carriers)	Black or Red	Gray or Roan	Black-some brindling	Black-can be spotted	Mostly Black
Red		Red	Red or Roan (cream if Charolais)	Red-some brindling	Red or Black, can be spotted	Red to mixed
White			White or Cream	White to gray	Spotted to white	Mixed
Light Hair, Dark skin				Light hair, dark skin to gray	Gray-can be spotted	Mixed
Spotted					Spotted	Mixed
Mixed Color						Mixed

respectively. There is another set of alleles that controls the dilution, or intensity, of that color. Dilution causes black to be muted to gray and red to be muted to yellow. As an example, Charolais cattle are red, but possess two alleles for dilution, which results in white coat color (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1994163/>). This is why Charolais x Angus cattle are gray (diluted black). A very thorough discussion of coat color in cattle, including its many variations (Table 3) can be found at (<http://simmental.org/site/pdf/other/olsoncolor.pdf>).

In a typical sale barn market, cattle are sold with little, if any, information made available about breed or performance. Many buyers will estimate performance (growth, carcass characteristics, etc.) in relation to the reputation of the breed; thus, they may look for signs that indicate a certain breed or breeds within crossbred cattle. Other buyers may be looking to source animals that qualify for black- or red-hided branded beef programs and are willing to pay a premium for these types of calves. Some breeds are prone to producing calves that have certain distinguished color markings, such as white-faces, brindling or white stockings on their legs.

Some general rules can be utilized to give the greatest chance of obtaining uniformly-colored groups of calves. Because red is recessive to black coat color, breeding solid red cows and bulls will produce solid red calves, which makes solid red an easy color to maintain in a crossbreeding system. However, because black is dominant to red, breeding solid black bulls and cows will often produce black calves, but may also yield red calves. To ensure a solid black calf crop, breed solid colored females (without diluter genes) to a homozygous black bull. If black baldy calves are desirable, use of Hereford bulls on black cows (or black bulls on Hereford cows) will yield the desired effect. If color extremes in the cowherd are a major concern, they can be masked by breeding through several generations to Charolais bulls.

In today's market, where marketing branded beef is advantageous, knowledge of coat color inheritance is essential. Even with traditional marketing systems, uniformity in coat color can often provide premiums/discounts in the market place. Therefore, knowing the basics of color inheritance will help in planning crossbreeding systems that create animals suitable for desired marketing niches or branded programs.

Table 3. Description of known coat color patterns in cattle (adapted from Olson 1999).

<i>Color or color modifier</i>	<i>Allele</i>	<i>Inheritance</i>	<i>Description</i>
Basic color (Extension)	Black (ED) Wild-type (E+) Red (e)	ED > E+ > e	Responsible for most coat color variation, wild-type is a brownish black sometimes observed in Brown Swiss, Jersey, Brahman, and Longhorn, other coat colors are modifications of these three basic colors
Brindle	Br	Dominant to no brindling	Brindling is alternating stripes of black and red pigmentation, animals must be homozygous wild-type to observe brindling
Agouti	Patterned blackish wild-type modifier (Apb) White-bellied modifier (aw) Fawn/dorsal stripe (ai)	Incompletely understood	These alleles are typically responsible for removal of either black or red color or both in specific areas of the body, such as along the underline or back
Dilution	Charolais (Dc) Simmental (Ds) Dun (Dn)	Charolais almost completely dominant, Simmental incompletely dominant to normal coloring	Charolais dilution is strong leading to light gray, cream, or white animals, Simmental dilution (also found in Gelbvieh, Longhorn and others) is moderate dilution of red and black, and dun is strong removal of red pigment and reduced removal of black pigment
Spotting	Hereford pattern (SH) Pinzgauer pattern (SP) Piebald (s)	S ^H => S ^P >S ⁺ >s	SH is Hereford pattern at five points when homozygous (white face only when heterozygous with non-spotting), Pinzgauer is variable white along topline and underline, piebald is irregular areas of pigment and extremities are usually white
Roan	R	Codominant with normal coloring	Homozygotes are almost completely white and heterozygotes are roan (red or black and white are codominantly expressed)
Belting	Bt	Dominant	White belting around midsection
Blaze	Bl		White face, often only a blaze when heterozygous
Brockling	Bc		Areas of pigmentation within areas of white spotting produced from other modifiers
Color-sided	Cs	Partially dominant	Homozygotes will have white body with pigmented ears, muzzle and feet (such as White Park) and heterozygotes show color-sided pattern with white dorsal stripe (can be roan) and roan on head

The Oklahoma Cooperative Extension Service

Bringing the University to You!

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
- It is administered by the land-grant university as designated by the state legislature through an Extension director.
- Extension programs are nonpolitical, objective, and research-based information.
- It provides practical, problem-oriented education for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.
- It utilizes research from university, government, and other sources to help people make their own decisions.
- More than a million volunteers help multiply the impact of the Extension professional staff.
- It dispenses no funds to the public.
- It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
- Local programs are developed and carried out in full recognition of national problems and goals.
- The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
- Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. 1214 GH