



# Recommended Vaccination Schedules for a Comprehensive Dairy Herd Health Program

Dan N. Waldner, Ph.D.  
Assistant Professor  
Extension Dairy Specialist

John Kirkpatrick, DVM  
Associate Professor  
Veterinary Clinical Sciences

Terry W. Lehenbauer, DVM  
Associate Professor  
Veterinary Pathobiology

Feeding practices, management styles, health care programs, and facilities vary greatly among dairy operations. Because of this, the degrees of stress, the patterns of disease resistance, and pathogen exposure are variable and unique to each operation. Consequently, there is no "one size fits all" vaccination program, but each program must be "tailor-made" to fit the individual needs of each dairy.

Immunization is a necessary aid to limit or prevent disease in cattle due to common agents, such as Infectious Bovine Rhinotracheitis (IBR) virus, Parainfluenza-3 (PI<sub>3</sub>) virus, Bovine Viral Diarrhea (BVD) virus, Bovine Respiratory Syncytial Virus (BRSV), clostridial infections, and leptospirosis. The design of a vaccination program must take into account a variety of factors including infectious disease problems in the immediate area or region. It is strongly recommended that producers contact a qualified veterinarian before proceeding with any herd health vaccination program.

## Killed (inactivated) vaccines (KV) vs Modified-live vaccines (MLV)

The most common vaccines on the market are either killed or modified-live. Most killed vaccines are provided in a liquid form that is ready for immediate use. Modified-live vaccines usually come as a dry powder that must be reconstituted prior to use. They should be used within a few hours after reconstitution, and they are particularly sensitive to harsh environmental conditions and should be protected from extreme temperatures or exposure to direct sunlight. Modified-live vaccines contain the disease organism that has been altered so that it may reproduce after it is injected into the host (animal); it is designed to develop immunity against the agent without making the animal sick. Modified-live vaccines are generally not recommended for pregnant animals. (See Table 1.) It takes approximately two to four weeks after the initial vaccination for the animal's body to develop sufficient protection against a challenge from the infectious agent included in the vaccine. Yearly boosters are usually recommended for both modified-live and killed vaccines. However, there are many new products on the market that are exceptions with regard to frequency of administration and handling as compared to older, conventional vaccines. It is important that all vaccines are handled and stored properly to maintain their potency, and

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: <http://osufacts.okstate.edu>

always read and follow the instructions carefully. Table 1 lists advantages and disadvantages of both vaccine types.

## Heifers (birth to 6 months of age)

Colostrum management is of primary importance for the health of the newborn calf. Calves should be fed 3 quarts of colostrum within an hour of birth and should receive a total of 6 quarts of colostrum within the first 24 hours of life.

Table 2 outlines the recommended vaccination schedule for dairy heifers from birth to six months of age. Young pre-weaned calves usually require little in the way of vaccines. In fact, a period of suppression or refractoriness to immunization may exist for two to three weeks after birth in calves that have received adequate transfer of antibodies from colostrum. However, an intra-nasal dose of IBR-PI<sub>3</sub> vaccine during this time can be justified as a preventative measure against respiratory disease related to those viral agents.

At weaning and two weeks prior to group penning or housing, calves should be vaccinated with a modified-live IBR, BVD, PI<sub>3</sub>, BRSV vaccine and a 7-way clostridial bacterin-toxoid. *Pasteurella spp. (multocida and haemolytica\*)* bacterin-toxoid and *Leptospira spp.* bacterin may be administered at this time.

At 6 to 10 months of age, heifers should be revaccinated with modified live IBR, PI<sub>3</sub>, BVD, BRSV virus vaccine, a minimum 2-way clostridial vaccine (up to 7-way), brucellosis vaccine, and leptospirosis bacterin (up to 5-way, pomona minimum). If a history of leptospirosis or clostridial disease exists on the property in calves at a younger age, then appropriate earlier vaccination should be done.

## Heifers (pre-breeding to calving)

Table 3 *bovis* bacterins within five days of mastitis vaccines. Animals should be vaccinated for Vibriosis (*Campylobacter fetus*) if using natural service.

## Adult cows

Table 4 outlines the recommended vaccination schedule for adult dairy cows. Animals must not receive any other gram negative vaccines including: *Pasteurella spp.*, *Salmonella spp.*, *Campylobacter sp.*, *Haemophilis somnus*, *E. coli* or *Moraxella bovis* bacterins within five days of mastitis vaccines. Annual vaccination for Vibriosis is recommended for non-AI herd. If

\* *Pasteurella haemolytica* has been renamed *Mannheimia haemo-*

a problem exists with scours, annual vaccination for Rota and Corona virus, *E coli*, and *Clostridium perfringens* (type C and D) may be considered to support other improved preventative management practices.

## Bulls

Table 5 outlines the recommended vaccination program for dairy herd bulls. It is suggested that producers purchase only virgin bulls from reliable sources with known vaccination histories, and these bulls should have negative test results for persistent BVD virus infection, brucellosis, and tuberculosis (depending on local and state requirements). All herd additions should also undergo a minimum isolation period of 30 days and preferably 60 days. All bulls should have been designated as satisfactory potential breeders, utilizing the standard breeding soundness examination, prior to exposure to breeding females.

## Summary

It is important that the owner and veterinarian develop a vaccination program tailored to fit the needs of the herd, and that it is implemented within current or proposed management practices. Pregnancy status of breeding females and herd/animal immune status are also important factors to be considered in any herd health program.

Vaccines, vaccine selection, and proper vaccine and equipment handling are only one component of a well-structured herd health program. If there is an absence of proper biosecurity (inter-herd) and biocontainment (intra-herd) practices, proper nutrition, and proper management of environmental conditions and other stress factors, any well-planned vaccination program may be ineffective. Always include a veterinarian in preventative herd health decisions, and always read and adhere to product instructions and precautions.

**Table 1.** Comparison of advantages and disadvantages of modified-live and killed (inactivated) vaccines.

Modified-live vaccines	Killed (inactivated) vaccines
<ul style="list-style-type: none"> <li>◆ Provide longer duration and more complete immunity</li> <li>◆ May produce better cellular and secretory immunity</li> <li>◆ Do not require multiple vaccinations for immunologic memory</li> <li>◆ Often do not require revaccinating or require fewer revaccinations during the life of an animal</li> <li>◆ Rarely causes hypersensitivities, but may be virulent to certain individual animals or may cause animals to produce clinical signs of disease</li> <li>◆ When used on pregnant animals some abortions may occur (depending on the particular viral agent)</li> </ul>	<ul style="list-style-type: none"> <li>◆ Provide short-lived systemic immunity</li> <li>◆ May provide limited cellular and secretory immunity</li> <li>◆ Often require revaccination to ensure immunologic memory</li> <li>◆ Require multiple vaccinations to maintain active immunity</li> <li>◆ Can cause hypersensitivity reactions</li> <li>◆ Cannot cause disease even in immunologically compromised animals</li> </ul>

**Table 2.** Recommended vaccination schedule for dairy heifers from birth to 6 months of age.

Age or time of administration	Disease	Type of vaccine or therapy
0-6 hours		Colostrum
6 weeks	IBR-PI <sub>3</sub> -BVD-BRSV Clostridial spp.	Modified live vaccine 7-way bacterin/toxoid
4-6 months	Brucellosis <sup>a</sup>	RB51
6 months	IBR-PI <sub>3</sub> -BVD-BRSV Clostridial spp. Leptospirosis	Modified live vaccine 7-way bacterin/toxoid 5-way bacterin

<sup>a</sup>Follow state and federal regulations: replacement heifers should receive immunization between 4 to 10 months at the owner's discretion depending on marketing strategies.

**Table 3.** Recommended vaccination schedule for heifers pre-breeding to calving.

Age or time of administration	Disease	Type of vaccine or therapy
Pre-breeding (10-12 months)	IBR-PI <sub>3</sub> -BVD-BRSV	Killed or modified live vaccine
	Leptospirosis	5-way bacterin
	<i>Clostridial spp.</i>	7-way bacterin/toxoid
	Vibriosis (optional)	Bacterin
40-60 days prior to calving	IBR-PI <sub>3</sub> -BVD-BRSV	Killed vaccine
	Leptospirosis	5-way bacterin
	Calf scours: Rota and Corona virus	Killed vaccine
	<i>E. coli + Clostridium perfringens</i> , type C & D	Bacterin/toxiod
3 weeks prior to calving	Calf scours: Rota and Corona virus	Killed vaccine
	<i>E. coli + Clostridium perfringens</i> , type C & D	Bacterin/toxiod
	Coliform mastitis	Bacterin

**Table 4.** Recommended vaccination schedule for adult dairy cattle.

Age or time of administration	Disease	Type of vaccine or therapy
40-60 days prior to calving	IBR-PI <sub>3</sub> -BVD-BRSV <sup>a</sup>	Killed vaccine
	Leptospirosis <sup>b</sup>	5-way bacterin
	Vibriosis (optional)	Bacterin
	Calf scours: Rota and Corona virus	Killed vaccine
	<i>E. coli + Clostridium perfringens</i> , type C & D	Bacterin/toxiod
3 weeks prior to calving	Calf scours: Rota and Corona virus	Killed vaccine
	<i>E. coli + Clostridium perfringens</i> , type C & D	Bacterin/toxiod
Follow label directions	Coliform mastitis	Bacterin

<sup>a</sup>Annual booster is necessary<sup>b</sup>Vaccination is recommended every six months in some areas.**Table 5.** Recommended vaccination schedule for dairy herd bulls.

Age or time of administration	Disease	Type of vaccine or therapy
Breeding soundness examination	IBR-PI <sub>3</sub> -BVD <sup>a</sup>	Killed vaccine
	Leptospirosis <sup>a</sup>	5-way bacterin
	Vibriosis <sup>a</sup>	Bacterin

<sup>a</sup>Annual booster is necessary

## The Oklahoma Cooperative Extension Service

### *Bringing the University to You!*

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
- It is administered by the land-grant university as designated by the state legislature through an Extension director.
- Extension programs are nonpolitical, objective, and research-based information.
- It provides practical, problem-oriented education for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.
- It utilizes research from university, government, and other sources to help people make their own decisions.
- More than a million volunteers help multiply the impact of the Extension professional staff.
- It dispenses no funds to the public.
- It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
- Local programs are developed and carried out in full recognition of national problems and goals.
- The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
- Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert E. Whitson, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. 0607