



Factors Affecting Wheat Germination and Emergence in Hot Soils

David Marburger
Small Grains Extension Specialist

One of the most cost-effective ways to increase fall forage production by winter wheat is early sowing. Sowing prior to September 20 generally provides enough time for wheat to establish canopy, produce some forage, and develop sufficient roots to anchor the plant in the soil. Therefore, dual-purpose wheat in the southern Great Plains is typically sown from late August through late September. Factors such as seed dormancy, high temperature germination sensitivity and coleoptile length may delay germination or prevent wheat seedling emergence when sown early into hot soils. Delayed germination reduces the amount of time for wheat growth prior to winter dormancy, resulting in less forage production. It is important for dual-purpose wheat producers to understand factors that might delay wheat germination or reduce emergence.

Post Harvest Dormancy

Wheat seed is considered dormant when it will not germinate, even when favorable conditions for germination are present. Some degree of seed dormancy is a favorable trait for a wheat variety. Seed dormancy is highest just before harvest. This dormancy can help prevent premature sprouting prior to harvest. Seed dormancy is gradually lost with time, but the rate of seed dormancy loss is affected by several factors. Inhibitory substances found in the seed coat of hard red winter wheat varieties, for example, can strengthen post-harvest dormancy. Wheat stored under extremely hot or cold conditions after harvest will generally germinate more readily than seed stored at ambient air temperatures. This is why it is recommended to place seed samples in the refrigerator for a day or two prior to running germination tests. The strength of dormancy also increases with decreasing temperatures during grain fill. So, seed of the same variety harvested from different areas of the state or region might behave differently when sown early. Post-harvest dormancy of most hard red winter wheat varieties will sufficiently dissipate by October. Early-sowing provides less time between harvest maturity and wheat sowing, and seed dormancy may not have dissipated enough to allow germination.

High Temperature Germination Sensitivity

In addition to post-harvest dormancy, some varieties have seed dormancy accentuated by high soil temperature, which is commonly referred to as high temperature germination sensitivity (Table 1). Wheat can germinate in soil temperatures from 40 F to 99 F, but temperatures from 54 F to 77 F are considered optimal. Soil temperatures in western Oklahoma

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: <http://osufacts.okstate.edu>

frequently exceed 80 F well into the fall. Soil temperatures at Altus, for example, typically exceed 85 F until mid-September and frequently do not fall below 80 F until October (Figure 1). These temperatures are sufficient to prolong dormancy in sensitive varieties such as 2174, Iba, LCS Mint, Overlay, Ruby Lee, SY Llano and WB-Grainfield. Varieties with high temperature germination sensitivity are generally not the best choice for sowing prior to September 15. A good rule of thumb

Table 1. High temperature germination sensitivity and earliest recommended planting dates for hard red winter wheat varieties.

<i>No sensitivity</i> <i>September 1</i>	<i>Moderate</i> <i>sensitivity</i> <i>September 10</i>	<i>Sensitive</i> <i>September 20</i>
Armour	Brawl CL Plus	2174
Avery	Byrd	Bentley
Billings	Deliver	Centerfield
Duster	Doans	Cutter
Endurance	Doublestop CL Plus	Iba
Gallagher	Everest	Jagalene
Greer	Fannin	Larry
Guymon	Fuller	LCS Mint
Jagger	Jackpot	LCS Wizard
Lonerider	Joe	Long Branch
OK Bullet	LCS Pistol	Overlay
Pete	NF 101	Ruby Lee
T154	OK Rising	Stardust
T158	Smith's Gold	SY Drifter
TAM 112	Spirit Rider	SY Flint
Winterhawk	T153	SY Grit
	WB-Cedar	SY Llano
		SY Monument
		TAM 204
		Tatanka
		WB4303
		WB4458
		WB4515
		WB4721
		WB-Grainfield
		WB-Redhawk
		Zenda




Figure 1. Average maximum observed temperature at a 4-inch depth under bare soil at Cherokee (top) and Altus, OK (bottom) from 2007 to 2016. Observed temperatures at shallower depths were likely higher. Data are from Oklahoma Mesonet.

is to plant less sensitive varieties, such as Duster, Winterhawk or Gallagher first and wait to sow sensitive varieties until after the soil temperature has cooled. In most years, the combination of time after harvest and cooler soil temperatures is sufficient to allow germination for most hard winter wheat varieties by September 15.

Coleoptile Length

Hot soil conditions at sowing also reduce coleoptile length. The coleoptile is a rigid, protective structure that covers the emerging shoot to aid in reaching the soil surface. Once the coleoptile breaks the soil surface, it stops growing and the first true leaf emerges. If the coleoptile does not emerge through the soil surface, the first true leaf emerges below ground, takes on an accordion-like appearance and the wheat plant typically dies (Figures 2A and 2B). For this reason, wheat should never be sown deeper than the coleoptile length. Wheat coleoptile length is related to mature plant height, and most



Figures 2A and 2B. If the wheat coleoptile fails to break the soil surface, the first true leaves will emerge below the soil surface. If this happens, the leaves will have an accordion-like appearance, as shown in the pictures, and the plants will die.

modern, semi-dwarf varieties have shorter coleoptiles than old, tall varieties. Most modern wheat varieties can safely be sown up to 1.5 inches deep, but most will not emerge when sown deeper than 1.5 inches into hot soils. For this reason, “dusting in” early-sown wheat and waiting on a cool rain to reduce high-temperature germination sensitivity and increase coleoptile length frequently results in more uniform emergence than planting deeper to reach moisture.

When sowing early, producers should carefully choose varieties and avoid those with varietal characteristics that can reduce germination. Reduced or erratic wheat germination will result in less fall forage production and reduced stocking rates. Soil temperatures are generally cool enough to allow full germination of most hard red winter wheat varieties by October 1.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: eeo@okstate.edu has been designated to handle inquiries regarding non-discrimination policies: Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU's Title IX Coordinator 405-744-9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President for Agricultural Programs and has been prepared and distributed at a cost of 20 cents per copy. Revised 0917 GH.